

Comparison between clinical estimated fetal weights (CEFW) Versus Ultrasonographic estimated fetal weight (UEFW) for co-relation with actual birth weight (ABW) in 3rd Trimester of PregnancyRabia Razaq^{1*}, Sajida Parveen², Muhammad Umer Razaq³, Sadaf Mubeen⁴, Fakhria Ijaz⁵, Shanza Ghaffar⁶¹Department of Gynecology and Obstetrics, Allied Hospital Faisalabad (38000), Pakistan²Department of Gynecology and Obstetrics, Lady Willington Hospital Lahore (54810), Pakistan³Department of Pediatric Medicine, Mayo Hospital Lahore (54810), Pakistan⁴Department of Gynecology and Obstetrics, Lady Aitcheson Hospital Lahore (54810), Pakistan⁵Department of Gynecology and Obstetrics, Lady Willington Hospital Lahore (54810), Pakistan⁶Department of Gynecology and Obstetrics, Lady Willington Hospital Lahore (54810), Pakistan**ARTICLE INFO***Article history:**Received: 28 December, 2019**Accepted: 21 January, 2020**Online: 14 March, 2020**Keywords:**Clinical**Actual Birth Weight**Fetal Weight**Ultrasonography**3rd Trimester**Pregnancy*

This article is open access under terms of Creative Commons Attribution License 4.0, which permits unrestricted use, distribution and reproduction in any medium provided the original work is cited properly.

DOI: 10.37978/gulf.v3i1.81**Correspondence:*****rabiarazaq@gmail.com*****ABSTRACT****Background:** Accurate prenatal estimation of birth weight is useful in the management of labour and delivery.**Objective:** To determine the correlation between clinical estimated fetal weight with actual birth weight in 3rd trimester of pregnancy and to determine the correlation between Ultrasonographic fetal weight assessment with actual birth weight in 3rd trimester of pregnancy.**Material & Methods:** This cross sectional study with non-probability purposive sampling technique was conducted in three tertiary care hospitals of Punjab, Department of Obstetrics & Gynaecology, Allied Hospital, Faisalabad, Lady Aitcheson Hospital Lahore and Lady Willington Hospital Lahore. Informed consent was obtained from each female to use their data for research purpose. Demographic details were also noted. Then females undergo CEFW was done by using Johnson's formula. Then ultrasonography was done on every female by experienced radiologists to get UEFW. FW measurement was done by using Shepard formula. Then females were followed-up till delivery of fetus. Actual birth weight (ABW) was noted on birth. Pearson correlation was used to measure the correlation coefficient for CEFW and UEFW with ABW. $P\text{-value} \leq 0.05$ was taken as significant.**Results:** In our study the mean age of the patients was 29.60 ± 6.23 years and the mean gestational age of 33.30 ± 2.31 weeks. The mean BMI value of the patients was 23.08 ± 1.26 Kg/m², the mean CEFW value 2219.60 ± 556.41 grams while the mean UEFW value of the patients was 2227.77 ± 521.94 grams and the mean value of ABW of the patients was 2284.00 ± 515.29 grams. In our study the positive correlation was found between the CEFW, UEFW with ABW of the baby.**Conclusion:** Our study results concluded that both the clinical estimation ultrasonography estimation showed the feasible and reliable results. Both showed positive correlation with actual birth weight.**Introduction**

Accurate prenatal estimation of birth weight is useful in the management of labour and delivery, permitting obstetricians to make decisions about instrumental vaginal delivery, trial of labour for patients suspected of having a low birth weight or macrosomic fetus (1). According to the existing literature, there is no truly accurate technique for evaluating FW (2).

Before delivery, accurate estimation of FW can have a major approach for decision and management of labour, perinatal outcome can be improved better (3). Estimation of FW can be done by external abdominal measurements, alone or associated with fundal height measurement and/or USG scan near 32 weeks

(4). Since the advent of ultrasound and its dissemination over the last three decades, and despite the lack of conclusive evidence, there has been a widespread belief that ultrasound is more accurate than other methods for predicting fetal weight (2).

A study correlate clinical estimation of fetal weight (CEFW) and Ultrasonographic estimation of fetal weight (UEFW) and estimated the correlation coefficient taking actual birth weight (ABW) as gold standard. Correlation between CEFW and ABW ($r = 0.074$) was insignificant and was almost showed no relationship while a significant correlation between UEFW and ABW ($r = 0.782$) (5). But another study reported that the correlation coefficient for the CEFW was 0.78 and UEFW was 0.74 and it was statistically demonstrated that both showed

significant positive correlation ($p < 0.001$) (6).

Rationale of this study is to compare the correlation between clinically estimated and actual birth weight and Ultrasonographic estimated and actual birth weight in 3rd trimester of pregnancy. Literature has reported that estimation of fetal weight during last trimester can be helpful in planning labour and delivery and achieving better maternal and perinatal outcome. But there is discrepancy in literature. Through this study we want to confirm that which method i.e. ultrasonography or clinical estimation of fetal weight is the best method to estimate FW before birth and can have a better fetal outcome in case of low FW. In routine, in tertiary care hospitals, obstetricians rely on ultrasonography but in sub urban areas or peripheries, facility of ultrasonography lacks. So we want to assess the reliability of FW clinically, so that we can rely on this method in future to lessen the burden and use of on USG. Objectives of study were to determine the correlation between clinical estimated fetal weights with actual birth weight in 3rd trimester of pregnancy and to determine the correlation between Ultrasonographic fetal weight assessment with actual birth weight in 3rd trimester of pregnancy.

Materials and Methods

This cross sectional study was conducted in three tertiary care hospitals of Punjab, Department of Obstetrics & Gynecology, Allied Hospital, Faisalabad, Lady Aitcheson hospital Lahore and Lady Willington Hospital Lahore from January 2019 to December 2019. Sample size of 100 cases was calculated with 5% type I error, 10% type II error and taking expected value of correlation coefficient i.e. 0.74 (6) between CEFW and ABW in 3rd trimester of pregnancy. Non-probability purposive sampling technique was used.

Ethical approval was taken from hospital ethical committee. Females of age 18-40 years with presenting in third trimester (gestational age of >30 weeks) with single cephalic fetus (on USG) with BMI 18-25 kg/m² were included in study. While patients with membrane rupture, multiple pregnancy, high risk patients like gestational diabetes (BSR > 200 gm/dl), PIH (BP > 140/90 mmHg), pre-eclampsia (PIH with protein urea +1 on dip stick) or eclampsia (pre-eclampsia with convulsions), anemia (Hb < 8 gm/dl), deranged LFTs (ALT > 40 IU, AST > 40 IU), deranged RFTs (serum Creatinine > 102 gm/dl) and those who had big uterine fibroid, polyhydramnios, oligohydramnios and congenital fetal anomaly (on USG) were excluded from study. Written informed consent was obtained from each female to use their data for research purpose. Demographic details (name, age, gestational age, BMI and parity) were also noted. Then females undergo CEFW was done by using Johnson's formula (as per operational definition). Then ultrasonography was done on every female by experienced radiologists to get UEFW. FW measurement was done by using Shepard formula (as per

operational definition). Then females were followed-up till delivery of fetus. Actual birth weight (ABW) was noted on birth (as per operational definition). All this information was recorded in the proforma. The data was entered and analyzed through SPSS version 20. Mean and SD were calculated for quantitative variables like age, gestational age, parity, CEFW, UEFW and ABW. Pearson correlation was used to measure the correlation coefficient for CEFW and UEFW with ABW. $P\text{-value} \leq 0.05$ was taken as significant.

Results

In this study total 100 patients participated. The mean age of the patients was 29.60 ± 6.23 years with minimum and maximum ages of 18 & 40 years respectively. The study results showed that the mean gestational age of the patients was 33.30 ± 2.31 weeks with minimum and maximum gestational ages of 30 & 37 weeks respectively. Frequency distribution of the patients showed that 23 (23%) patients appeared with no parity, the patients with parity one was 26 (26%), patients with parity two were 30 (30%), patients with parity three were 13 (13%) and 8 (8%) patients appeared with parity four. In this study the mean height of the patients was 168.19 ± 7.033 cm with minimum and maximum heights of 150 & 177 cm respectively. In this study the mean weight of the patients was 65.32 ± 5.33 kg with minimum and maximum weights of 52 & 75 kg respectively.

Table 1: Descriptive statistics of study population

	Age	Gestational age	Height	Weight in Kg	BMI in Kg/m ²
n	100	100	100	100	100
Mean	29.60	33.30	168.19	65.32	23.08
SD	6.23	2.31	7.033	5.33	1.26

The study results showed that the mean BMI value of the patients was 23.08 ± 1.26 Kg/m² with minimum and maximum BMI values of 20.10 & 25 Kg/m² respectively. The study results showed that the mean SFH value of the patients was 26.32 ± 3.58 cm with minimum and maximum SFH values of 21 & 34 cm respectively. The study results showed that the mean CEFW value of the patients was 2219.60 ± 556.41 grams with minimum and maximum CEFW of 1395 & 3410 grams respectively. In this study the mean BPD value of the patients was 83.75 ± 5.90 grams with minimum and maximum BPD values of 74 & 94 grams respectively.

Table 2: Descriptive statistics of fetus

	AC gram	UEFW (grams)	Actual birth weight (grams)	SFH (cm)	CEFW in grams	BPD in gram
n	100	100	100	100	100	100
Mean	292.7	2227.77	2284.00	26.3	2219.6	83.75
SD	22.25	521.94	515.29	3.58	556.41	5.90




Figure 1: Frequency Distribution of Parity

In this study the mean AC values of the patients was 292.74 ± 22.25 grams with minimum and maximum AC values of 255 & 330 grams respectively. The study results showed that the mean value of UEFW of the patients was 2227.77 ± 521.94 grams with minimum and maximum UEFW values of 1465 & 3177.40 grams respectively. The study results showed that the mean value of ABW of the patients was 2284.00 ± 515.29 grams with minimum and maximum ABW values of 1400 & 3400 grams respectively. In our study the positive correlation was found between the clinical estimated fetal weight and the actual birth weight of the baby. i.e $r=0.965$. Our study results showed the positive correlation between the ultrasonographic estimated fetal weight and the actual birth weight of the baby. i.e $r=0.927$.

Discussion

This present cross sectional study was conducted at Unit I, Department of Obstetrics & Gynecology, Allied Hospital, Faisalabad to determine the correlation between clinical estimation of fetal weight and ultrasound estimation of fetal weight with actual birth weight.




Figure 2: Correlation between CEFW & ABW




Figure 3: Correlation between UEFW & ABW

The screening and management of abnormal fetal growth, whether it macrosomia or growth restriction, remain important objectives of prenatal care. In a low-risk and unselected population, such screening is based mainly on a series of ultrasound examinations. Estimating fetal weight is an easy and straight forward way in which to monitor fetal growth and to screen for intrauterine growth restriction (IUGR) (7).

SFH measurement is a simple and inexpensive method to detect abnormal fetal growth; however, according to a recent systematic review, there is not enough evidence to evaluate the use of this technique in the routine antenatal care (8). In our study the mean value of SFH of the patients was 26.32 ± 3.58 cm. Alessandra Curt et al (9) showed in their study that the role of obstetric and maternal factors in birth weight prediction at term of pregnancy is confirmed. The value of the variables used to build up the statistical algorithm is higher to clinical estimation performed in labor by an expert obstetrician (10-13).

Our study results showed that the both clinical and ultrasound estimation of fetal birth are positively correlated with actual birth weight. In our study the mean CEFW value of the patients was 2219.60 ± 556.41 grams and it has positive correlation with ABW ($r=0.965$), similarly the mean value of UEFW of the patients was 2227.77 ± 521.94 grams and it had positive correlation with ABW ($r=0.927$).

NilgünGüdücü et al (14) concluded in their study that ultrasonographic fetal weight estimations correlate with the actual birth weight better when performed in the late third trimester, but ultrasonographic fetal weight estimation early in the third trimester may allow for better follow up and planning of delivery both in small and large for gestational age fetuses.

Akinola S. Shittu et al demonstrated in their study that the accuracy of clinical estimation was highest in the birthweight range of 2,500–<4,000 g and lowest for the low-birthweight group (<2,500 g). This is in consonance with what several investigators have shown that the clinical method is best for

estimating fetal weight in the reference birth-weight range of 2,500 to <4,000 g with accuracy (mean absolute percentage error) of ± 7.5 –19.8% depending on gestational age and that below 2,500 g (15-21).

Peregrine et al (22) concluded in their study that clinical estimates of birth weight perform favorably compared with ultrasonographic estimates, ultrasound immediately prior to labor is more accurate at predicting the low- or high-birth-weight fetus.

One more study showed that Clinical estimation of birth weight in early labor is as accurate as routine ultrasonic estimation obtained in the preceding week. In the lower range of birth weight (less than 2500 g), ultrasonic estimation is more accurate; in the 2500–4000 g range, clinical estimation is more accurate. In the higher range of birth weight (greater than 4000 g), Both methods have similar accuracy (23). Correlation between CEFW and ABW ($r = 0.074$) was insignificant and was almost showed no relationship while a significant correlation between UEFW and ABW ($r = 0.782$) (5). But another study reported that the correlation coefficient for the CEFW was 0.78 and UEFW was 0.74 and it was statistically demonstrated that both showed significant positive correlation ($p < 0.001$) (6). Ben-Haroushet al(104) explains that there was a high correlation between EFW and birth weight ($R(2) = 0.775$, $P < 0.001$). The mean birth weight was 3207 \pm 561 g, and mean absolute weight difference was 227 \pm 197 g; (absolute range, 0-1700 g; actual range, - 986 to + 1700 g).

Sanyal P et al (105) also showed a good correlation between the ultrasound measurements and the postnatal measurements i.e. $r^2 = 0.98$. Akinola S et al (15) described that correlation coefficient for ultrasound estimation (0.74) Uotila et al. in their comparison of ultrasonic estimation showed (0.77) correlation. Akinola S et al (15) described the correlation coefficient of clinical estimation (0.78) while Dare et al. also showed the similar proportion (0.74).

Conclusion

Our study results concluded that both the clinical estimation and ultrasonography estimation showed the feasible and reliable results. Both showed positive correlation with actual birth weight. So in future we can rely on clinical method in areas where ultrasound facility is not available.

Conflict of Interest: This study has no conflict of interest to declare by any author.

Disclosure: None

Human and Animal Rights: No rights violated.

References

- Manzanares, S., Gonzalez-Escudero, A., Gonzalez-Peran, E., López-Criado, M., & Pineda, A. (2019). Influence of maternal obesity on the accuracy of ultrasonography birth weight prediction. *The Journal of Maternal-Fetal & Neonatal Medicine*, 1-6: 10.1080/14767058.2019.1567708.
- Preyer, O., Husslein, H., Concin, N., Ridder, A., Musielak, M., Pfeifer, C., & Husslein, P. (2019). Fetal weight estimation at term—ultrasound versus clinical examination with Leopold's manoeuvres: a prospective blinded observational study. *BMC pregnancy and childbirth*, 19(1), 122: <https://doi.org/10.1186/s12884-019-2251-5>
- Kong, C. W., & To, W. W. K. (2019). Comparison of the accuracy of INTERGROWTH-21 formula with other ultrasound formulae in fetal weight estimation. *Taiwanese Journal of Obstetrics and Gynecology*, 58(2), 273-277: 10.1016/j.tjog.2019.01.019.
- Liao, K., Tang, L., Peng, C., Chen, L., Chen, R., Huang, L., & Chen, C. (2019). A modified model can improve the accuracy of foetal weight estimation by magnetic resonance imaging. *European journal of radiology*, 110, 242-248.
- Sharma, K. A., Das, D., Dadhwal, V., Deka, D., Singhal, S., & Vanamail, P. (2019). Two dimensional fetal biometry versus three-dimensional fractional thigh volume for ultrasonographic prediction of birthweight. *International Journal of Gynecology & Obstetrics*, 145(1), 47-53.
- Pushpamala Ramaiah, D., Elsayed, L. A., Lindsey, G., & Johargy, A. (2019). Estimation of Fetal Size and Weight using Various Formulas: [10.31142/jtsrd23231](https://doi.org/10.31142/jtsrd23231)
- Kadji, C., Cannie, M. M., Resta, S., Guez, D., Abi-Khalil, F., De Angelis, R., & Jani, J. C. (2019). Magnetic resonance imaging for prenatal estimation of birthweight in pregnancy: review of available data, techniques, and future perspectives. *American journal of obstetrics and gynecology*, 220(5), 428-439: 10.1016/j.ajog.2018.12.031
- Janas, P., Radoń-Pokracka, M., Nowak, M., Staroń, A., Wilczyńska, G., Brzozowska, M., & Huras, H. (2019). Effect of oligohydramnios on the accuracy of sonographic foetal weight estimation in at term pregnancies. *Taiwanese Journal of Obstetrics and Gynecology*, 58(2), 278-281.
- Al-Shawwa, M. O., & Abu-Naser, S. S. (2019). Predicting Birth Weight Using Artificial Neural Network.
- Abbas, H., Badar, S., Javed, Z. and Ramadan, M.A.A. 2020. Level of Serum Uric Acid in Pre-eclamptic and Normal Pregnant Women. *Int J Front Sci*, 4(1), 20-22: 10.5281/zenodo.3613298
- Fong, Z. V., Pitt, H. A., Strasberg, S. M., Molina, R. L., Perez, N. P., Kelleher, C. M., ... & Chang, D. C. (2019). Cholecystectomy during the third trimester of pregnancy: proceed or delay? *Journal of the American College of Surgeons*, 228(4), 494-502: 10.1016/j.jamcollsurg.2018.12.024
- Ross, K. M., Cole, S. W., Carroll, J. E., & Schetter, C. D. (2019). Elevated pro-inflammatory gene expression in the third trimester of pregnancy in mothers who experienced stressful life events. *Brain, behavior, and immunity*, 76, 97-103: 10.1016/j.bbi.2018.11.009.
- Sanderson, M., Kostoula, M., Palazzo, F., Boret, T., & Kong, C. (2019, November). A case of successful parathyroidectomy in the third trimester of pregnancy. In *Society for Endocrinology BES 2019* (Vol. 65). BioScientifica.
- Faure-Bardon, V., Magny, J. F., Parodi, M., Couderc, S., Garcia, P., Maillotte, A. M., & Pladys, P. (2019). Sequelae of congenital cytomegalovirus following maternal primary infections are limited to those acquired in the first trimester of pregnancy. *Clinical infectious diseases*, 69(9), 1526-1532: 10.1093/cid/ciy1128.
- Panwar, M., Kumari, A., Arora, R., Singh, V., & Banswal, R. (2019). Raised neutrophil lymphocyte ratio and serum beta hCG level in early second trimester of pregnancy as predictors for development and severity of preeclampsia. *Drug discoveries & therapeutics*, 13(1), 34-37: 10.5582/ddt.2019.01006.